

Deliverable 24: Documentation and database for TREHS software tool

April 2017

Contact:
F.E.M. Research Group,
Departament d'Ecologia, Facultat de Biologia, UB
Diagonal, 643 - 08028 Barcelona +34 93 403 11 88

lifetrivers@ub.edu
www.lifetrivers.eu



LIFE13 ENV/ES/000341

With support from the LIFE+ programme of the European Union



Duration: 06/2014 – 06/2018

Implementing the Water Framework Directive to temporary rivers: tools for the assessment of their ecological status
LIFE13 ENV/ES/000341

Project partners:



Authors: Francesc Gallart (IDAEA- CSIC), Jérôme Latron (IDAEA- CSIC), Pilar Llorens (IDAEA- CSIC), Núria Cid (UB-FEM), Núria Bonada (UB-FEM), Narcís Prat (UB-FEM), Antoni Munné (ACA), Carolina Solà (ACA), Mònica Bardina (ACA), Teodoro Estrela (CHJ), Arancha Fidalgo (CHJ), Sara-Maria Jiménez-Argudo (CHJ), Rosa-María Vega (CHJ), Justin Jeuffroy (ENGEES), Antoni Hernández-Casahuga

Contributors: Andreu Manzano (ACA), Javier Sanromán (CHE) and Rogelio Galván (CHE).

Deliverable 24: Documentation and database for TREHS software tool
Responsible beneficiary: IDAEA-CSIC

Design by: © GLOBAL DESIGN DEVELOPMENT, S.L.

CONTENT

ABSTRACT	4
1 INTRODUCTION	5
2 HYDROLOGICAL FEATURES OF THE TREHS TOOL	6
3 PRINCIPLES USED IN TREHS	7
3.1. Spatial and temporal units	7
3.2. Aquatic states and aquatic phases	7
4 DATA GATHERING AND MANAGEMENT	8
4.1 Flow discharges	8
4.2. Interviews	9
4.3 In situ observations and aerial or ground-level photographs	9
5 DATA ANALYSES AND OUTPUTS	10
5.1 Temporary regime metrics	10
5.2 Temporary regime graphs	10
5.3 River regime terminology and classification	12
5.4 Assessing hydrological alteration.....	13
5.4.1 <i>Reference regime conditions</i>	14
5.4.2 <i>Hydrological status</i>	14
6 APPLICATION OF TREHS IN ECOLOGICAL STATUS ASSESSMENTS	15
7 DISCUSSION AND PERSPECTIVES	15
7.1 Design changes during the development and interim application of TREHS.....	15
7.2 Weaknesses of TREHS	16
7.3 Perspectives.....	17
8 ACKNOWLEDGEMENTS	17
9 REFERENCES	17
APPENDIX A: GRAPHS USED IN TREHS	21




Abstract

When the regime of a river is not perennial, there are four main difficulties with the use of hydrographs for assessing hydrological alteration: i) the main hydrological features relevant for biological communities are not quantitative (discharges) but qualitative (phases such as flowing water, stagnant pools or lack of surface water), ii) stream flow records do not inform on the temporal occurrence of stagnant pools, iii) as most of the temporary streams are ungauged, their regime has to be evaluated by alternative methods such as remote sensing or citizen science, and iv) the biological quality assessment of the ecological status of a temporary stream must follow a sampling schedule and references adapted to the flow- pool-dry regime.

To overcome these challenges within an operational approach, the freely available software tool TREHS has been developed within the EU LIFE TRIVERS project. This software permits the input of information from flow simulations obtained with any rainfall-runoff model (to set an unimpacted reference stream regime) and compares this with the information obtained from flow gauging records (if available) and interviews with local people, as well as instantaneous observations by individuals and interpretation of ground-level or aerial photographs.

Up to six metrics defining the permanence of water flow, the presence of stagnant pools and their temporal patterns of occurrence are used to determine natural and observed river regimes and to assess the degree of hydrological alteration. A new regime classification specifically designed for temporary rivers was developed using the metrics that measure the relative permanence of the three main phases: flow, disconnected pools and dry stream bed.

Finally, the software characterizes the differences between the natural and actual regimes, diagnoses the hydrological status (degree of hydrological alteration), assesses the significance and robustness of the diagnosis and recommends the best periods for biological quality samplings.



1 Introduction

Despite being largely ignored in the past, rivers that recurrently cease to flow at some point in time and/or space or may dry out completely (hereafter termed temporary, following Uys and O’Keeffe, 1997) are probably the most common fluvial ecosystems in the world (Fritz et al., 2006; Nadeau and Rains, 2007; Datry et al., 2014). Recent decades have seen still more of them, due to climate change and pressures on water use, such as water abstraction (Jacobsen, 2004; Larned et al., 2010). Flow regime shifts from perennial to temporary are predicted for many regions in the world under future climate change scenarios (Kirkby et al., 2011; Döll & Schmied, 2012). In fact, temporary rivers are already frequent in Mediterranean basins; for example, over 20% of river water bodies in Spain’s Júcar River Basin are temporary (CHJ, 2016).

In consequence, significant basic and applied research on temporary rivers has increased since the late 1980’s (Sheldon, 2005; Leigh et al., 2016b). Much of this research has focused on understanding biodiversity patterns and trends, especially because global change is increasing flow temporariness in many parts of the world and threatening local biodiversity (Thoms & Sheldon, 2002; Blanchette & Pearson, 2012; García-Roger et al., 2011). Although hydrological and ecological assessment issues currently occupy much of the research into temporary rivers undertaken in many countries (Leigh et al., 2016b), the lack of information and of new developments in hydrology are the main impediments to advancing in the science and management of temporary rivers (Acuña et al., 2004; Leigh et al., 2016b; Seaman et al., 2016; Skoulikidis et al., 2017).

Ecological assessment of rivers has a long tradition in freshwater ecology (e.g. Bonada et al., 2006; Friberg et al., 2011). In most countries, ecological status is assessed by using biological quality indices to compare biological communities sampled at a site with its reference (benchmark or baseline) conditions (Hawkins et al., 2010). However, as reference conditions may vary over time due to wet or dry periods (Munne & Prat, 2011), changes in river regime should also be taken into account. In Europe, the Water Framework Directive (WFD, European Commission, 2000) requires that assessment of ecological status must also include information on the hydromorphological alterations that support biological elements, such as the hydrological regime, river continuity or the geomorphological conditions. However, this is still a challenging task in temporary rivers (Reyjol et al., 2014; Skoulikidis et al., 2017) because appropriate classification and assessment of the hydrological regime are required before it can be decided whether the current methods developed for perennial systems are still valid or new approaches need to be developed.

Hydrological alterations due to human activity are now one of the main causes of impairment of riverine ecosystems locally and globally. Determining them is the first step in river restoration and conservation (International River Foundation, 2007; Poff et al., 2010; European Commission, 2012; Seaman et al., 2016). Thus, distinguishing whether a water body has a natural or artificial hydrological regime is vital for the setting of environmental objectives and adequate assessment of ecological status (Skoulikidis et al., 2017; Stubbington et al., 2017a). In the case of temporary rivers, a proper assessment of these alterations is of even greater relevance (Leigh et al., 2016a; Cid et al., 2017). For instance, perennial rivers can become temporary due to water abstractions, but in some particular cases, rivers that are naturally temporary may become artificially perennial, usually as a result of effluent inputs from waste-water treatment plants (Luthy et al., 2015). For example, this occurs in the intensively exploited aquifers of the Vinalopó river in Spain, whose aquifer water levels have decreased in recent years by between 65 m and 350 m due to groundwater abstractions, leading to the river drying (Custodio et al., 2016). Current streamflows are due mainly to effluent inputs from waste-water treatment plants (CHJ, 2016).

In perennial rivers, several software tools, such as IHA (Indicators of Hydrologic Alteration; Richter et al., 1996), IHARIS (Indicators of Hydrologic Alteration in RIVERs; Martínez Santa-María and Fernández Yuste, 2010) or DRIFT (Downstream Response to Imposed Flow Transformations; Brown et al., 2013), are currently used to determine flow regimes and hydrological alterations by comparing impacted against reference (unimpacted) hydrographs. However, these tools usually fail when applied to temporary rivers because i) the main hydrological features relevant to biological communities in temporary rivers are not quantitative (i.e. discharges), but qualitative (i.e. the presence of flowing water, stagnant pools or the lack of surface water) (Boulton, 1989, 2000; Uys and O’Keeffe, 1979; Buffagni et al., 2009; Seaman et al., 2016); ii) river flow records do not report the temporal occurrence of stagnant pools (Gallart et al., 2016), which act as refuges for many species during the cessation of flow (e.g. Bonada et al., 2006; Buffagni et al., 2009; Stubbington et al., 2017a); iii) as most temporary rivers are ungauged, hydrographs are often unavailable and those derived from models are of doubtful validity (Seaman et al., 2016); iv) those temporary rivers with a gauging station usually have only old data, which do not allow analysis of the current hydrological regime.

Furthermore, although the conceptual bases for identifying hydrological regimes of temporary rivers are established (Uys and O’Keeffe, 1997), in practice there are myriad terminologies and classifications based only on flow statistics. These classifications are usually difficult to implement or have unproven ecological significance (Poff and Ward, 1989; Poff, 1996; Uys and O’Keeffe, 1997; Rossouw et al., 2005; Mackay et al., 2012; Gallart et al., 2012; Arthington et al., 2014; Datry et al., 2016; Skoulikidis et al., 2017), to such an extent that classifications do not allow extrapolation of

observations and do not help the comparison of regimes with reference conditions (Seaman et al., 2016). Thus, as temporary rivers are of great importance in some regions, particularly in the Mediterranean, Water Management Administrations in the European Union need specific tools for the implementation of the WFD in these cases.

Within this conceptual and methodological context, and with the explicit objective of implementing the WFD for temporary rivers, the hydrological part of the Temporary Rivers Ecological and Hydrological Status (TREHS) free software tool was developed. Although it was conceived for operational use, this does not prevent its being a useful research tool, since it includes some cutting-edge concepts and permits the gathering of information that is not compulsory under the current requirements of the WFD, but may help in the understanding of riverine ecosystems and contribute to their sound management.

The main aim of this work is to propose updated methods for the operational monitoring, assessment and classification of the regime of temporary rivers, along with the evaluation of the degree of hydrological alteration. These targets are preconditions to setting environmental objectives and applying appropriate biomonitoring protocols in this type of water body. These methods are assembled in the TREHS software tool which makes the gathering, storing, analysis and evaluation of the information easier. A second but more ambitious aim is to propose the methods described below as a blueprint for monitoring and cataloguing temporary river regimes wherever they occur.

This is not, however, a TREHS user guide or a substitute for it. This guide can be downloaded at <http://www.lifetrivers.eu>. Examples of application of TREHS to real rivers are shown in Deliverable 25 of this LIFE TRIVERS project.

2 Hydrological features of the TREHS tool

The hydrological features of TREHS, in particular the classification of regimes and the assessment of the degree of hydrological alteration, were conceived as an instrument for the investigation of temporary river regimes, not focusing on water resources but rather on the ecological implications relevant to their knowledge and management.

TREHS is a user-friendly interface and a data base management instrument for the gathering of diverse types of hydrological-related data of temporary rivers. It calculates six temporary regime metrics from these data, offers a classification of the river regime on the basis of three metrics and has a graphical interface that facilitates river regime visualisation, classification and comparison between diverse river sites. In addition, if reference conditions are provided, TREHS assesses the hydrological regime (i.e. Hydrological Status), based on the evaluation of the degree of hydrological alteration. TREHS has the following functions, reported in a sequential order:

- Description of the test reach (station) with the adequate meta-data; identification of the operator, date, location, River Basin District, water body and local name. The tool makes it possible to use interactive maps for the location of the reaches.
- Collection from different data sources: there are three kinds of data, flow records, interviews and observations. Flow records can be obtained from gauging stations or model-simulated flow series; diverse time periods of flow records or simulations from different models can be separately analysed. Observations may consist of in situ observations from the field and observations from aerial or ground-level photographs.
- Calculation of six temporary river metrics and their variability, using all the diverse data sources stated above.
- Analysis of the long-term monthly or seasonal relative frequencies of the aquatic states of the water body by drawing the Aquatic States Frequency Graph. This function can also be obtained from various data sources.
- Drawing two types of graphs for showing and comparing the metrics results obtained from the diverse data for a single river reach or a group of them.
- Classification of the river aquatic phases regime, taking into account the metrics that represent the relative occurrence of the main three aquatic phases: flow, disconnected pools and dry channel.
- Advice on the best sampling calendar to be used for collecting samples and assessing the Ecological Status of the water body by including the hydrological constraints of aquatic life.
- Comparison between the actual and the reference (non-impacted) river regime metrics for assessing the degree of hydrological alteration in terms of the implications for the development of macroinvertebrate aquatic fauna. Determination of Hydrological Status as a measurement or diagnosis of hydrological alteration due to human pressures on river regime (water quantity).
- Listing metrics and reporting results for a selected river reach or a group of them.

3 Principles used in TREHS

TREHS follows the conceptual framework of hydrological alteration and its ecological limits (Richter et al., 1996; Poff et al., 2010; Seaman et al., 2016) and the description of the hydrological phases in temporary rivers (Boulton, 2003; Fritz et al., 2006). TREHS also implements the concepts and applications for investigating and managing temporary rivers formulated during the European Community's Seventh Framework Programme "Mediterranean Intermittent River Management" (MIRAGE) research project (Gallart et al., 2012; Prat et al., 2014; de Girolamo et al., 2015; Cid et al., 2016; Nikolaidis et al., 2013) and uses alternative data-gathering methods (Turner & Richter, 2011; Buytaert et al., 2014; Datry et al., 2016; Gallart et al., 2016). Table 1 shows the main acronyms and terms used in TREHS that were defined in Gallart et al. (2016) or are newly defined in this work.

In particular, TREHS is based on the working hypothesis that the hydrological regime features that are most relevant to the development of aquatic life in a temporary river are not the temporal patterns of water discharges but of the occurrence of mesohabitats (e.g. Larned et al., 2010; Gallart et al., 2012; Schriever et al., 2015). This is consistent with the idea that temporary rivers should be seen as a distinct class of systems, instead of just hydrologically challenged permanent rivers (Larned et al., 2010).

All questions of data gathering, analysis and evaluation turn on these premises, with the advantage that qualitative states are much more easily obtained through non-instrumental methods than water discharges are (Fritz et al., 2006; Turner and Richter, 2011; Datry et al., 2016; Gallart et al., 2016).

3.1. Spatial and temporal units

The spatial unit for the management of river systems under the regulations of the WFD is the water body (European Commission, 2000), which may comprise several kilometre-long river sections where regime may spatially vary. This is why TREHS software focuses mainly on a part of a water body, a hundred meter-long river reach called "station". Several stations may be defined and separately analysed by TREHS within each water body according to their spatial regime variation or data availability. Results are not mapped in TREHS, but it provides suitable outcomes that can be implemented in a GIS.

TREHS uses several temporal units. The basic time unit used is the month, as community composition of many aquatic organisms change among months and recovery from water quantity disturbances (i.e. floods and drying periods) usually occur in few weeks (Robinson et al., 2003; Leigh et al., 2016a; Dolédec et al., 2017). Nevertheless, when interviews or either in situ or photographic observations are used, the month becomes a too short unit because it is not feasible to get sufficiently time-detailed information; in these cases, for practical reasons, the season is taken as the operational time unit. The metrics described below in section 5.1 use diverse time units, from the month to the semester, whereas the year is the time period used to complete the river regime. Some issues arisen with the use of time units, such as the role of short events and the loss of information when flows are aggregated from a daily scale to a monthly one are analysed in the discussion (section 7).

3.2. Aquatic states and aquatic phases

The units for describing the immediate hydrological state at the moment of inspection are the aquatic states, defined as the transient sets of aquatic mesohabitats occurring on a given river reach at a particular moment, depending on the hydrological conditions (Gallart et al., 2012). Six aquatic states were there defined, from wet to dry: flooding conditions (Hyperrheic), full prevalence of all the possible mesohabitats (Eurheic), sequence of pools connected by flowing water threads (Oligorheic), occurrence of isolated pools (Arheic), disappearance of surface water, with the wet alluvium still allowing underground aquatic life (Hyporheic) and the desiccation of the river bed and alluvium, involving the disappearance of any active aquatic habitat (Edaphic).

The aquatic states occurring at the moment of sampling are identified for better evaluation of the biological assemblages (Prat et al., 2014) and are recorded in the TREHS data base. Nevertheless, as it is not currently possible to obtain this detailed information for the past, to obtain the metrics and classification of the river regime the six aquatic states have been simplified to three aquatic phases (flowing water, isolated pools and dry river bed). The identification of the temporal patterns of occurrence of these three phases is consistent with the conceptual 'continuum' defining the behaviour of temporary river regimes (Uys and O'Keeffe, 1997), may be successfully obtained with various non-instrumental methods and characterizes the river regime sufficiently for most purposes (Datry et al., 2016; Gallart et al., 2016). The relationships between the three aquatic phases and the biological communities using communities as predictors of hydrological conditions have been explored in Cid et al. (2016).

4 Data gathering and management

Input to TREHS may come from three data sources: i) flow discharge sequences (measured or simulated), ii) interviews and iii) in situ observations from field visits or through aerial or ground-level photographs. These data are stored and analysed by the software in order to obtain different outputs (see section 5).

4.1 Flow discharges

Monthly flow data from a gauging station or model simulations are used to obtain the statistics of the occurrence of diverse aquatic states, following the method described in Gallart et al. (2012, 2016). This kind of information allows only the appropriate determination of the aquatic states that correspond to the flowing water phase (Hyperrheic, Eurheic and Oligorheic), separated from the remaining ones that correspond to zero flows, once flow thresholds between these states are assessed. To identify these thresholds correctly, field observations on the aquatic states synchronous with discharge measurements would be needed. However, in the absence of these observations, thresholds can be provisionally assessed with the help of the shape of the flow duration curve (distribution function of flow discharges), as proposed by TREHS.

Accurate determination of the discharge reading corresponding to real zero flow is usually not straightforward for several reasons: gauging stations are not normally designed to measure zero flows and small changes due to erosion/deposition of sediments in the gauging section or inadequate maintenance of the artificial control can modify the zero flow value. In addition, some gauging stations may not measure water flow through the alluvium, whereas other stations are designed to intercept subsurface flow within the alluvium in order to measure it as surface flow. Therefore, low flow values may correspond to either the Arheic (isolated pools) or Hyporheic (no surface water) states (an example is shown in Deliverable 25). Furthermore, low flow values on the monthly scale may actually mean a month with a few days with flow among many days without flow, but with or without pools.

When flow simulations obtained with a rainfall-runoff model are used, it must not be forgotten that, apart from the role of the above issues in the flow data used for model calibration, most current models are not designed to simulate zero water discharges.

In practice, the user can decide the value corresponding to actual zero flow and can assign the smaller values to either the Arheic or Hyporheic states. TREHS states a warning message that the frequency of the Arheic state (pools phase) is usually underestimated by flow measurements.

Several flow records of diverse origin or period and selected thresholds between aquatic states may be stored in TREHS and alternatively selected to compare results. The reference (unimpacted) conditions are usually taken from flow values simulated with a rainfall-runoff model assuming natural conditions, but flow records measured in the past before hydrological alteration in the same or another location can also be used.

Table 1: Acronyms, terms and concepts used in the paper. ¹: defined in this work; ²: defined in Gallart et al., 2012.

ACRONYM	CONCEPT	DEFINITION
AF	Alternate-Fluent ¹	<u>Aquatic phases regime</u> from FPD plot; alternation between the three aquatic phases, but flowing for longer periods ($Mf > 0.40$ and $Md \geq 0.10$).
AI	Alternate ¹	<u>Aquatic phases regime</u> from FPD plot; alternation between the three aquatic phases ($Mf \leq 0.40$, $Mp < 0.40$ and $Md < 0.60$).
Aquatic phase	Aquatic phase ¹	Simplified transient stage of temporary water condition occurring in a reach and moment (flow, standing pools, dry stream bed)
Aquatic state	Aquatic state ²	Transient set of mesohabitats occurring in a reach and moment, controlled by water stage (<i>Hyperrheic</i> , <i>Eurheic</i> , <i>Oligorheic</i> , <i>Arheic</i> , <i>Hyporheic</i> , <i>Edaphic</i>)
Arheic	<i>Arheic</i> ²	<u>Aquatic state</u> ; zero surface flow but isolated water pools are present.
AS	Alternate-Stagnant ¹	<u>Aquatic phases regime</u> from FPD plot; alternates between the three aquatic phases, but remains stagnant for longer periods ($Mf \leq 0.40$, $Mp \geq 0.40$ and $Md \geq 0.10$).
ASFG	Aquatic States Frequency Graph ²	<u>River regime graph</u> ; relative frequency of the aquatic states during months or seasons throughout the year.
Edaphic	<i>Edaphic</i> ²	<u>Aquatic state</u> ; waterless river bed and alluvium, involving the disappearance of any active aquatic habitat.
Ep	Episodic ¹	<u>Aquatic phases regime</u> from FPD plot; usually dry river with either flowing or stagnant water at infrequent intervals ($Md \geq 0.80$).
ESs	Equinox-solstice seasonality ¹	<u>Metric</u> ; temporal arrangement of no flow periods: the relative frequency of 0-flow months in equinoxes minus the one in solstices.
Eurheic	<i>Eurheic</i> ²	<u>Aquatic state</u> ; water discharge is high enough to allow the occurrence and connectivity of all the feasible aquatic habitats in the reach.
FPD	Flow-Pool-Dry plot ¹	<u>Aquatic phases regime graph</u> ; triangle showing the complementary metrics on

ACRONYM	CONCEPT	DEFINITION
FS	Fluent-Stagnant ¹	the permanence of the three aquatic phases in the river reach (<i>Mf</i> , <i>Mp</i> and <i>Md</i>). <u>Aquatic phases regime</u> from FPD plot; usually flowing but otherwise with isolated pools ($0.40 < Mf < 0.90$ and $Md < 0.10$).
Hyperrheic	<i>Hyperrheic</i> ²	<u>Aquatic state</u> ; infrequent high water (flood) conditions
Hyporheic	<i>Hyporheic</i> ²	<u>Aquatic state</u> ; most of the stream bed is devoid of surface water in the reach, although alluvium may remain wet enough to allow hyporheic life.
Md	Dry channel permanence ¹	<u>Metric</u> ; long-term mean annual relative number of months without surface water in the channel.
Mf	Flow permanence ²	<u>Metric</u> ; long-term mean annual relative number of months with flowing water.
Mp	Pools permanence ¹	<u>Metric</u> ; long-term mean annual relative number of months with isolated pools.
Oc	Occasional ¹	<u>Aquatic phases regime</u> from FPD plot; river usually dry that sometimes, but not often, has flowing or stagnant water ($0.60 \leq Md < 0.80$).
Oligorheic	<i>Oligorheic</i> ²	<u>Aquatic state</u> ; surface water discharge is scarce but sufficient to connect most pools in the reach through water threads.
Pe	Perennial ¹	<u>Aquatic phases regime</u> from FPD plot; permanently flowing ($Mf > 0.99$).
Qp	Quasi-Perennial ¹	<u>Aquatic phases regime</u> from FPD plot; usually flowing, except on infrequent occasions ($0.90 < Mf \leq 0.99$).
Sd₆	Six-month predictability of zero-flow periods ²	<u>Metric</u> ; temporal arrangement of no flow periods: the unity minus the relative frequency of the zero-flow months in the wetter 6-month period divided by the relative frequency in the complementary (drier) 6-month period.
St	Stagnant ¹	<u>Aquatic phases regime</u> from FPD plot; river usually in the form of isolated water pools ($Mf \leq 0.40$ and $Md < 0.10$).
SWs	Summer-winter seasonality ¹	<u>Metric</u> ; temporal arrangement of no flow periods: the relative frequency of 0-flow months in summer minus the one in winter.
TRP	Temporary Regime Plot ²	<u>Temporary regime graph</u> ; X-Y plot of flow permanence <i>Mf</i> (X) versus Six-month predictability of zero-flow periods <i>Sd₆</i> (Y).
WDF	Water Framework Directive	<u>Water regulation</u> : European framework for water policy (European Commission, 2000)

4.2. Interviews

Interviews are designed to assess the current regime of the river reach in the last 10 years period. They follow the method described in Gallart et al. (2016) and are designed to be answered by neighbours of the fluvial system, i.e. people living near the rivers or working in vegetable gardens beside the river, selected haphazardly from people met near the river and willing to answer the questions. Alternatively, interviews can be conducted with key informants who have a more expert professional or leisure relationship with the river regime.

Given the impracticality of obtaining information by means of these methods for each month and for all the aquatic states described above, the months are grouped in seasons and the six aquatic states are reduced to the three aquatic phases. Thus, after some preliminary questions, the core of the questionnaire consists of a template with four columns corresponding to the four seasons and three rows corresponding to three aquatic phases (flow, disconnected pools and dry river bed). As each box represents the number of months of occurrence of the corresponding state in the corresponding season, every column must cover 3 months and every row 12 months.

Interviewees are also asked about the occurrence of wet/saturated alluvium after the disappearance of surface water (pools). Yet, for the driest rivers, when the interviewee indicates that the river flow ‘ceases for long periods’, the question is reformulated in reverse as ‘How many days per year does the river carry water?’. This question was included in order to document low-flow frequency.

Any interview may be selected or discarded for the analysis, depending on the user’s confidence in its reliability.

4.3 In situ observations and aerial or ground-level photographs

The primary purpose of this data source input to TREHS is the documentation of the aquatic state of the reach at the date when it is visited for water and biological sampling. The entry was also adapted to include interpretations of past aerial photographs or surface photographs such as those taken by the Google Maps Street View multitemporal facility. For the last cases, as it is not possible to identify the six aquatic states in photographs, these have to be simplified into the three aquatic phases. In the current version of TREHS, the data from both in situ and photographic observations are merged for obtaining metrics and graphs, so the recorded aquatic states are simplified to aquatic phases when analysed.

Flow and pool frequency statistics are calculated from observations only when at least five observations are available and then, as a measurement of dispersion, the resolution of the statistics is calculated as the inverse of the number of observations. As in the case of interviews, the temporal scale for observations is simplified to the season scale. Statistics

and graphs on the temporal pattern of features are derived only when there are at least three observations per season. In this case, permanence statistics are first calculated for each season and subsequently averaged, in order to prevent any bias due to different numbers of observations in the various seasons. This prevention cannot be made when there are less than three observations per season, so in these cases the resulting permanence statistics may be subject to some seasonal bias.

Aerial or ground-level photographs may be taken at some distance from the station point, in which cases operators are advised to state the coordinates of the observation point. The capture date of aerial photographs may be unknown, as many orthophotographs are composed from photograph mosaics without a specific capture date; in these cases the seasonal information is lost and only the capture year may be recorded.

5 Data analyses and outputs

5.1 Temporary regime metrics

Six metrics, exclusive to temporary rivers and obtainable with the above-mentioned methods, were defined. Three metrics describe the time-compressed occurrence of the corresponding three aquatic phases, whereas the other three metrics try to describe the temporal arrangement of these phases. We decided not to use metrics on the six aquatic states because there is not yet sufficient information. The metrics selected are as follows:

Mf; flow permanence, defined as the long-term mean annual relative number of months with flow, with values between 0 (always dry) and 1 (always flowing, i.e. perennial river systems). This metric was defined in Gallart et al. (2012) from former studies (e.g. Poff, 1996 and Arscott et al., 2010).

Mp; pool permanence, defined as the long-term mean annual relative number of months when isolated pools occur, with values between 0 (pools never occur in the system) and 1 (always with pools, i.e. perennial still-water/lentic systems).

Md; dry channel permanence, defined as the long-term mean annual relative number of months when the channel has no surface water. It is a metric complementary to the two metrics above, so $Md=1-(Mf+Mp)$.

Sd₆; six-month predictability of zero-flow periods, defined in Gallart et al. (2012) and calculated by Eq. 1:

$$Sd_6 = 1 - \left(\frac{\sum_1^6 Fd_i}{\sum_1^6 Fd_j} \right) \quad (\text{Eq. 1})$$

where Fdi represents the multi-annual frequencies of 0-flow months for the contiguous 6 wetter months of the year and Fdj represents the multi-annual frequencies of 0-flow months for the remaining 6 drier months. Wet and dry 6-month periods mean here those with fewer and more zero-flow frequencies in the long term, respectively. This variable is dimensionless and takes the value of 0 when zero flows occur equally throughout the year in the long run and 1 when all the zero flows occur in the same 6-month period every year. When the regime is fully permanent, this metric cannot be computed, so the value of 1 is set to indicate full predictability.

SWs; summer-winter seasonality, defined as the difference in the relative frequencies of 0-flow months between summer and winter. It takes a value of 1 when there is no flow during summer versus continuous flow in winter and -1 when the contrary occurs. Summer and winter are calculated in TREHS as for the Northern Hemisphere; this metric would take the contrary sign in the Southern Hemisphere.

ESs; equinox-solstice seasonality, defined as the difference in the relative frequencies of 0-flow months between equinoxes and solstices. It takes a value of 1 when there is no flow during equinoxes versus continuous flow in solstices and -1 when the contrary occurs.

5.2 Temporary regime graphs

Three graphs can be obtained by TREHS: the Aquatic States Frequency Graph (ASFG) and Temporary Regime Plot (TRP), both defined in Gallart et al. (2012), and the new Flow-Pool-Dry plot (FPD).

The ASFG summarizes the relative frequency of the wetter aquatic states throughout the year, using a monthly temporal scale when it employs flow records or simulations or the three aquatic phases with a seasonal temporal scale when it employs interviews or observations (see Fig. A.1 in the Appendix A). The purpose of this graph is to show the relative importance of the diverse states or phases throughout the year and the degree of seasonality of the regime at a glance. It also gives a first impression of how the sampling calendar should be defined in the station. Nevertheless, it does not allow the quantitative assessment of the river regime required for comparisons between rivers or reaches.

The TRP (see Fig. A.2 in the Appendix A) was designed to compare the two main metrics relevant to the occurrence of flow obtained for diverse rivers. Thus, flow permanence (Mf) and seasonal (Sd6) predictability are represented in this plot. The grey triangle represents an area where the values of the two metrics are incompatible and the bars represent the standard error of the metrics. This plot can be used to compare the regimes of diverse rivers or the metrics obtained for the same river when diverse sources of information are used. Four sectors in this plot represent the temporary aquatic regime types defined by Gallart et al. (2012): Perennial (P), Intermittent-pools (I-P), Intermittent-dry (I-P) and Ephemeral (E). These types are shown in this graph for comparison but not used below, because a more ambitious classification was developed for TREHS on the basis of the plot subsequently described. The main drawback of the TRP plot shown above is the lack of information on the occurrence of surface water in the form of stagnant pools when flow is interrupted.

As both interviews and observations may provide relevant information on the frequency of pools that cannot be plotted in the TRP graph, a flow-pools-dry (FPD) plot was designed in order to show the metrics associated with the three aquatic phases relevant to aquatic life development (flow, disconnected pools and dry river bed) and observable in the river reaches with these methods. The design of the plot is triangular, the classic format when three complementary components are analysed, such as for sand, silt and clay components in soil texture (Soil Survey Division Staff, 1993). In the FPD plot (Fig. 1 and Fig. A.3 in the Appendix A), the axis on the left represents flow permanence (Mf), with the percentage of Mf increasing from the triangle bottom (Mf=0%) to the top (Mf=100%). The axis on the right represents pool permanence (Mp), increasing from the left axis (Mp=0%) to the bottom right vertex (Mp=100%). Finally, the axis on the bottom represents dry channel permanence (Md), complementary to the others ($Md=1-(Mf+Mp)$), with the lowest value in the right axis (Md=0%) and the highest value in the bottom left vertex (Md=100%).

Therefore, the upper, right and left vertexes of the FPD plot represent perennial riverine systems (i.e. perennial rivers), perennial lentic systems (i.e. perennial ponds or wetlands) and terrestrial systems, respectively. Points along the lower axis, with Mf=0% and different values of Mp and Md, represent temporary ponds or wetlands. Points along the left axis represent temporary rivers without a pool phase, whereas points on the right axis represent temporary rivers without any dry phase. Finally, the inside of the triangle represents temporary rivers that alternate between the three phases, with a wide range of different Mf, Mp and Md values (see section 5.1 for details).




Fig. 1: Arrangement of the three main metrics that correspond to the three aquatic phases; flow permanence (Mf), Isolated pools permanence (Mp) and dry river permanence (Md), in the FPD (Flow – Pools – Dry) graph. The arrows show the progression of every one of the three metrics whereas the axes show the values of every one of them. The central point represents a river that undergoes the three aquatic phases with the same frequency.

5.3 River regime terminology and classification

The five metrics described in section 5.1 and the graphs shown in 5.2 may be used to characterize and compare the regimes of diverse rivers, but an operational and ecologically relevant classification of the regimes was considered necessary for clearer communication in this multidisciplinary field (Uys and O’Keeffe, 1997), extrapolation of observations and progress in the sound management of rivers (Poff et al., 2010; Seaman et al., 2016). As argued in the Introduction, there is consensus among authors that aquatic life in temporary rivers depends not only on the occurrence of flow but also on the presence of surface water in the form of stagnant pools when flow is interrupted (e.g. Robson et al. 2013; Davis et al. 2013). Some pools may persist through months of no rainfall whilst others may change in size or disappear, for reasons difficult to be identified (Seaman et al., 2016). Consequently, there are some terminologies and classifications of the regime of temporary rivers that mention the occurrence of pools, but fail to operationally include their frequency in the identification of regime classes due to the lack of adequate statistics (e.g. Uys and O’Keeffe, 1997; Rossouw et al., 2005; Gallart et al., 2012).

The new aquatic phases regime classification implemented in TREHS was designed as i) fully applicable from available information, ii) taking into account the statistics of the three aquatic phases, iii) able to be represented in a single graph, iv) conflict-free from the most usual terminologies, and v) defined from hydrological features assumed to have biological implications, though these are not yet proved. However, practical reasons made it appropriate to discard the representation of the temporal structure of the aquatic phases. Therefore, it is to be expected that the biological significance of the classes designed will have different biological implications in distinct climate settings. Moreover, since both the terminology and the classification are coded in a spreadsheet auxiliary to the TREHS application, they can be updated by an advanced user using different threshold values for the metrics or even adding other TREHS metrics to the procedure. The approach selected for regime classification was based on the FPD plot (Fig. 2), using the following attributes:

- Perennial: Permanently flowing, except on rare occasions. The term ‘temporary’, following Uys and O’Keeffe (1997), is used as a blanket opposite term for the remaining rivers that occasionally or recurrently cease to flow.
- Fluent: Usually flowing.
- Stagnant: Usually takes the form of isolated pools.
- Alternate: Rotates between the three aquatic phases.
- Occasional: River usually dry that sometimes, but not often, has flowing or stagnant water.
- Episodic: Dry river with either flowing or stagnant water at infrequent intervals.

These terms are combined to identify nine types of regime, as shown in Table 2 and Fig. 2, where the threshold values defined for the three metrics are indicated. Two of these boundaries are assumed as the most relevant for aquatic life: Mf smaller than 0.4 is assumed as a practical boundary where usual WFD methods cannot be used to assess biological status; and Md smaller than 0.1 represents conditions with quasi-perennial surface water, either flowing or stagnant.

The results of this classification are displayed in TREHS for all the diverse types of information. Furthermore, as some European Member States included classifications of the rivers according to their natural flow regimes in the respective transpositions of the WFD (“ORDEN ARM/2656/2008” in Spain and “DECRETO 16 giugno 2008, n. 131” in Italy), these classifications are also offered by TREHS for the natural (reference) conditions, as the ‘prescribed regime’, as well as for each and every kind of information available.




Fig. 2: Distribution of the TREHS aquatic phases regimes in the FPD plot. Qp: Quasi-perennial; AF: Alternate-Fluent; FS: Fluent-Stagnant; St: Stagnant; AS: Alternate-Stagnant; AI: Alternate; Oc: Occasional; EP: Episodic. Mf: flow permanence; Mp: pool permanence; Md: dry channel permanence. The orange dots represent river stations where the metrics were obtained from in situ or photographic observations (73 points).

Table 2: Nomenclature and metrics boundaries of aquatic phases regimes as used in TREHS. Mf: flow permanence, Mp: pool permanence; Md: dry channel permanence. The characteristic metric boundaries used for defining the regimes in Fig. 2 are shown in **bold**.

Regime	Mf	Mp	Md
Perennial (Pe)	$0.99 < Mf \leq 1.00$	$0.00 \leq Mp < 0.01$	$0.00 \leq Md < 0.01$
Quasi-perennial (Qp)	$0.90 < Mf \leq 0.99$	$0.00 \leq Mp \leq 0.10$	$0.00 \leq Md \leq 0.10$
Fluent-Stagnant (FS)	$0.40 < Mf \leq 0.90$	$0.00 \leq Mp < 0.60$	$0.00 \leq Md < 0.10$
Alternate-Fluent (AF)	$0.40 < Mf \leq 0.90$	$0.00 \leq Mp < 0.50$	$0.10 \leq Md < 0.60$
Stagnant (St)	$0.00 < Mf \leq 0.40$	$0.50 \leq Mp < 1.00$	$0.00 \leq Md < 0.10$
characterAlternate-Stagnant (AS)	$0.00 < Mf \leq 0.40$	$0.40 \leq Mp < 0.90$	$0.10 \leq Md < 0.60$
Alternate (AI)	$0.00 < Mf \leq 0.40$	$0.00 \leq Mp < 0.40$	$0.20 \leq Md < 0.60$
Occasional (Oc)	$0.00 < Mf \leq 0.40$	$0.00 \leq Mp < 0.40$	$0.60 \leq Md < 0.80$
Episodic (Ep)	$0.00 < Mf \leq 0.20$	$0.00 \leq Mp < 0.20$	$0.80 \leq Md < 1.00$

5.4 Assessing hydrological alteration

After Article 4(1) of the WFD, the regime of a fluvial water body must be assessed for its ability to reach a good status of biological indicators (European Commission, 2000; CIS, 2015). Unfortunately, the ecology of temporary rivers is not yet sufficiently understood to make a sound evaluation of the ecological implications of regime alterations. Indeed, following the recommendations of the Ecological Limits of Hydrologic Alteration (ELOHA) framework (Poff et al., 2010), it is necessary i) to find ‘baseline’ or reference unimpacted regime characteristics for the water body under study,

ii) to classify the river regime using ecologically relevant variables, iii) to determine the deviation of the current regime from the baseline-condition one and iv) to develop regime alteration-ecological response relationships.

TREHS was designed to help cope operationally with the first three steps of the ELOHA framework and to determine Hydrological Status (HS) as an assessment of the ecological relevance of hydrological alteration, on the basis of expert criteria that can be easily updated when new information is made available. In the following subsections, the way the metrics described in section 5.1 are used for assessing the degree of hydrological alteration and determining its ecological relevance (HS) is summarized.

5.4.1 Reference regime conditions

The default method for determining the baseline or reference regime for a station in TREHS is the use of a flow-discharge series simulated with a rainfall-runoff model, assuming natural conditions. This kind of simulation for every water body is usually available for the implementation of the WFD (for instance, in Spain according to the “Instrucción de Planificación Hidrológica”, ORDEN ARM/2656/2008). Alternatively, historical flow records from gauging stations obtained before hydrological alteration can also be used to determine the reference regimes.

In these cases, simulated or recorded flow series are handled as described in section 4.1 to obtain the corresponding metrics and are selected as the reference ones. It is important to remember that the occurrence of isolated pools and therefore the determination of the Mp metrics are not straightforward when water flow data are used for calculating the metrics. The permanence of pools Mp obtained with these data is expected to be usually underestimated.

If reference regime metrics can be obtained from any other source (flow records from another gauging station in the area, interviews for this purpose), the corresponding metrics can be directly entered into TREHS. Finally, if diverse sources of reference regimes are available, such as from different rainfall-runoff models, their use in TREHS can be switched on and off in order to compare the results.

5.4.2 Hydrological status

TREHS calculates the degree of hydrological alteration with an expert scoring method from the differences between the metrics obtained for the reference regime and the actual one. These calculations are made on a separate auxiliary spreadsheet that can be inspected by the user in order to monitor the process and, if need be, update some of the expert criteria. To avoid a long description, an example of this spreadsheet is offered as supplementary material to this paper.

First, for every metric, the average and standard deviation of the values obtained from the diverse types of information on the reference and actual regime are obtained. The TREHS user can switch off any of the types of information if bias is suspected.

Then, the differences between the reference and the available actual metrics are compared with threshold values that depend on the reference Mf value, to decide whether the divergences are unacceptable; the more permanent the regime is, the lower divergences of the metrics are permitted. These threshold values are calculated in the auxiliary spreadsheet from tables of benchmark values that can be updated by an expert user.

The criteria used for assessing the hydrological alteration are as follows:

- Decrease of flow permanence Mf, two levels of severity (gentle and harsh).
- Decrease of surface water permanence (Mf+Mp), two levels of severity.
- Increase of flow permanence Mf, two levels of severity.
- Change of seasonal predictability Sd6.
- Change of seasonal patterns SWs or ESs.

Note that an increase in flow permanence or a change in the temporal pattern is also taken as hydrological alteration because they may facilitate the advent of generalist invasive species, particularly fishes and crayfishes (e.g. Riley et al., 2005).

Every criterion is penalised with one negative score that is subtracted from a value of 4; then the HS is determined as ‘not altered’, ‘lightly altered’, ‘moderately altered’ or ‘highly altered’ for total values from 4 to 1 respectively.

TREHS also displays the criteria used for this determination, in order to inform the manager of the measures to be taken for regime reclamation. Finally, it also calculates the degree of confidence of the diagnosis issued, based on the ratio between the metric differences and their standard deviations, as well as its robustness, based on the number of different kinds of information used.

6 Application of TREHS in ecological status assessments

The hydrological information provided by TREHS is intended to guide managers on the establishment of environmental objectives and selection of the most appropriate methods for ecological status assessment of temporary rivers.

On the one hand, hydrological status (i.e. distinguishing whether a water body has a natural or altered hydrological regime) contributes to defining specific environmental targets according to the river's natural regime and thereby to proposing adequate restoration or conservation measures. The WFD requires assessment of the ecological status of a river by means of groups of aquatic organisms, namely biological quality elements (i.e. macroinvertebrates, algae, macrophytes and fish). The combination of several indicators provides contrasting but complementary information on how they might respond differently to each stressor (Mykrä et al., 2012). Each taxonomic group is affected in a different way by regime alteration because each one has different traits as ways of coping with drying or floods, etc. For example, fish communities have limited traits to cope with interruption of flow or drying riverbeds (Kerezy et al., 2017), whereas macroinvertebrates and even algae have more resistant and resilient strategies (Sabater et al., 2017; Stubbington et al., 2017b). Thus, alteration of natural regimes that imply habitat loss, such as disconnected pools during the scarce or zero-flow period (e.g. alteration from a natural FS to Oc due to water abstraction), lead to stronger implications for the maintenance of native fish communities, which negatively affect the achievement of ecological status. In such examples, in which poor hydrological status is evident, biomonitoring of macroinvertebrate communities would not ensure a correct ecological status assessment and fish communities should be included (Benejam et al., 2010). Suitable knowledge of the natural presence of fish communities for each river site is essential prior to ecological status assessment with this biological quality element. Thus, TRESH software can provide useful information on the natural river regime that can later be compared with measured data, which shows whether fish fauna is expected to be present at each river site analysed.

On the other hand, river regime classification with information on permanence of flow (Mf), pools (Mp) and dry channel (Md), together with seasonal predictability (Sd6), may help managers to decide whether ecological status can be assessed by means of standard biomonitoring methods, mainly developed for perennial rivers, and thus partially avoid the exclusion of many temporary rivers from biomonitoring programs. For example, in a river with a flow permanence of 60-70% (SF or AF regime, see Table 2) showing high seasonal predictability, current methods could be used if the sampling calendar is adapted to the period with the greatest probability of finding lotic conditions (Prat et al., 2014). This is the case of many Mediterranean-climate rivers, in which the dry season is highly predictable (Bonada & Resh, 2013; Tonkin et al., 2017) and biotic indices based on macroinvertebrates have proved successful (Mazor et al., 2014; Munné & Prat, 2011; Cid et al. 2016).

However, even this adaptive approach could be limited in these types of rivers, depending on the level of aquatic habitat fragmentation (e.g. spatial isolation) during the dry season, which could lower organisms' recolonization potential (Datry et al., 2014b). In contrast, for a river with a flow permanence (Mf) and pool permanence (Mp) < 0.4, and with low seasonal predictability (e.g. Oc or Ep regimes, Table 2), current methods for the assessment of the ecological status are usually non-applicable. Even in reference sites, aquatic communities subjected to these regimes are usually poor in species and those that persist are tolerant to disturbances and human impacts; this will result in the underestimation of the ecological status and thus constrain the applicability of commonly used methods (Bonada et al., 2007; Buffagni et al., 2009; Cid et al., 2017; Munné and Prat, 2011). In these cases, characterized by the fact that most of the time rivers are dry and water flows sporadically in episodes of storms, its status should be evaluated preferably according to a hydromorphological assessment. Besides, the development of novel methods based on terrestrial communities can be a good solution (Sánchez-Montoya et al., 2016; Corti and Datry, 2015). Similarly, in those systems where pool habitats predominate throughout the year, specific methods could also be adapted and/or developed.

Further development of the TRESH software will include the methods to provide a complete evaluation for the ecological status of temporary streams.

7 Discussion and Perspectives

7.1 Design changes during the development and interim application of TREHS

TREHS was developed under the LIFE+ TRivers project and devised as a tool for applying the methods formulated during the MIRAGE project to the implementation of the WFD in temporary rivers. During its development and interim application, however, the original idea changed significantly, mainly due to the incorporation of information alternative to water discharge records, as well as the development of the FPD plot and regime classification with the advent of information on the occurrence of stagnant pools. The development of the FPD plot and the associated classification of regimes needed several months of discussion, but was established as a useful framework for deliberation, comparisons and communication.

TREHS was applied to 119 stations from the Catalan river basin district (ACA), the Júcar river basin district (CHJ) and the Ebro river basin district (CHE). The stations investigated were selected from stations in the first two districts where temporariness made the implementation of the WFD difficult, and from stations with zero-flow records in the third district. Alongside other methods, river regimes were assessed with the help of observations in 73 of these stations, as shown in Fig. 2. The most striking result is that these regimes cover most of the FPD plot, with at least two stations in every one of the regime classes. Two examples of TREHS application to real rivers are shown in Deliverable 25.

The hydrological part of TREHS has proved an advantageous tool for gathering, managing and analysing hydrological data from temporary rivers, particularly when there are no available flow records. Furthermore, it liberates researchers and managers from their traditional dependence on flow records or tentative model simulations, which may become disproportionate in these environments, while it offers new possibilities for a range of alternative observations, such as citizen science. At present, a smartphone application (Riu.neT) is being developed for making it easier and more functional to gather information on temporary river regimes and assess their ecological quality when possible.

In addition, the assessment of the regime of the temporary rivers investigated, as well as the ‘aquatic states’ background, became an essential framework for sound sampling and evaluation of the biological communities, as well as for comparing biological determinations between diverse investigated rivers.

7.2 Weaknesses of TREHS

Nevertheless, there are several weak points in TREHS that will need (if feasible) further development:

- The hydrological part of TREHS manages somewhat fuzzy information, mainly at the single station scale. This uncertainty is partly due to the intrinsic variability of the regime of temporary rivers in both time and space, and partly due to the constraints caused by limited sampling in the case of observations and by interviewee subjectivity in the case of interviews. Nevertheless, this information is much better than the frequent lack of flow records; and the user is advised on the levels of uncertainty and robustness associated with the data. He/she can, therefore, look for more information at the same station or at a nearby one, if necessary, in order to obtain the data necessary for an adequate hydrological assessment and diagnosis.

- In its present form, TREHS is not able to handle hydrological events occurring at shorter time scales than the month, whereas it is well known that runoff events triggered by storms are usually flashy and may occur in days or hours. As already stated in section 4.1, there are difficulties when daily flows are aggregated to the monthly scale. In fact, it is possible to store in situ or photographic observations of aquatic states at a time scale down to daily step in TREHS, but the analysis or display of this information is being made at the seasonal scale, looking for long-term patterns instead for a given time period. Shorter time steps may be used for analysing the recent history of aquatic states before and during biological sampling using the same principles (qualitative states instead of flow measurements, see an example in Gallart et al., 2012), but this cannot be made within TREHS.

- The metrics frontier values used to separate the regime classes in the FPD plot in Fig. 2 were tentatively defined in order to provide users with a clear terminology for cataloguing, comparing and exchanging the information gathered with interviews and observations. It may nevertheless happen that, when more data are available, the classes defined turn out to be excessively provisional and insufficient for assessing the hydrological controls on aquatic life. Further developments should aim at improving the classification, if sufficient biological data become available. Actually, TREHS uses the differences between the reference and actual regime metrics for determining the hydrological status, but not the respective regime classes defined in the FPD plot.

- As the TREHS classification of the regime of temporary rivers does not take into account the representation of the temporal structure of the aquatic phases, it may be expected that the biological significance of the classes designed will have different biological implications in diverging climate settings. This is not, however, an impediment to its use, because, if necessary, other climate categories can be added to the simpler flow-pools-dry classification (see an example in Uys and O’Keeffe, 1997).

- Flow discharge records, whether measured or simulated, were confirmed as an inadequate source of data for assessing the regime of temporary rivers. Not only is the measurement of water flow inadequate for correctly determining the occurrence of stagnant pools, but the design and inadequate maintenance of the gauging sections mean that there may be significant errors in the detection of low and zero flows. These limitations are particularly relevant when flows simulated with the help of a rainfall-runoff model are used to reproduce the natural flow regime, because they add the limitations associated with any hydrological modelling. Nevertheless, TREHS is not the cause of these limitations, but, on the contrary, is a useful tool for comparing diverse modelling approaches, not in terms of the simulated hydrographs, but in terms of the frequency and timing of zero-flow periods.

- TREHS is designed to be used at the station (reach) scale, while temporariness is also a spatial issue because diverse regimes may coexist at the same time along a sufficiently long river section. Given its current design, the analysis of

spatial patterns of temporariness must be handled point by point, and any management with a GIS must use the TREHS results off-line.

TREHS does not afford any useful approach for the quality assessment of rivers with Occasional or Episodic regimes. It only helps to define them in this class. Ongoing methods based on terrestrial organisms may be a valuable approach for improving these regimes' ecological assessment and management in the absence of sufficient aquatic characterisation (Sánchez-Montoya et al., 2016; Corti and Datry, 2015, Stubbington et al., 2017a).

7.3 Perspectives

The authors frankly expect that the developments gathered in TREHS will be a useful aid in the investigation and management of temporary rivers. In particular, the systematization of the collection and interpretation of data alternative to flow records, along with the development of the TREHS aquatic phases regime classification, may help to break through the confines imposed by the outdated approach of thinking of temporary rivers as simply hydrologically challenged perennial rivers instead of as a distinct class of ecosystems (Larned et al., 2010). Note that perennial rivers occupy just one point in the wide diversity of the FPD (Fig. 2) plot!

Indeed, these systematisations may provide the kernel of a useful conceptual and operational framework for spreading the study and monitoring of temporary rivers wherever they occur, such as the efforts being made in the SMIRES COST Action CA15113 (<http://www.smires.eu/>).

New data collecting methods, conceptual frameworks, metrics, terminology and classification are needed for this purpose. They should be discussed in collaborative international arenas such as the above-mentioned COST Action.

Meanwhile, we indeed recommend noting down the concurrent aquatic state of the temporary river reach when biological samples are taken.

8 Acknowledgements

This work is dedicated to the memory of the late Maria Rieradevall, a cheerful and committed colleague for many years and at the earlier stages of this project.

Aerial photographs were obtained from the Cartographic and Geological Institute of Catalonia (ICGC), the Instituto Geográfico Nacional (IGN) and Google Earth. Ground-level photographs were obtained from Google Maps' Street View. The spreadsheet for analysing three complementary variables developed by Graham and Midgley (2000) was used in the development of TREHS. We are also indebted to the interviewees, and to M. EAUDE for reviewing the English.

9 References

- Acuña V., Datry T., Marshall J., Barceló D., Dahm C.N., Ginebreda C.A., McGregor G., Sabater S., Tockner K. and Palmer M.A., Why should we care about temporary waterways?, *Science* **343**, 2014, 1080–1081.
- Arcott D.B., Larned S., Scarsbrook M.R. and Lambert P., Aquatic invertebrate community structure along an intermittence gradient: Selwyn River, New Zealand, *J. North. Am. Benth. Soc.* **29**, 2010, 530-545.
- Arthington A.H., Bernardo J.M. and Ilhéu M., Temporary rivers: linking ecohydrology, ecological quality and reconciliation ecology, *River Res. Appl.* **30**, 2014, 1209–1215.
- Benejam L., Angermeier P.L., Munné A. and García-Berthou E., Assessing effects of water abstraction on fish assemblages in Mediterranean streams, *Freshw. Biol.* **55**, 2010, 628–642, <http://dx.doi.org/10.1111/j.1365-2427.2009.02299.x>.
- Blanchette M.L. and Pearson R.G., Macroinvertebrate assemblages in rivers of the Australian dry tropics are highly variable, *Freshwater Sci.* **31**, 2012, 865–881.
- Bonada N. and Resh V.H., Mediterranean-climate streams and rivers: geographically separated but ecologically comparable freshwater systems, *Hydrobiologia* **719**, 2013, 1–29, <http://dx.doi.org/10.1007/s10750-013-1634-2>.
- Bonada N., Prat N., Resh V.H. and Statzner B., Developments in aquatic insect biomonitoring: a comparative analysis of recent approaches, *Annu. Rev. Entomol.* **51**, 2006, 495–523.
- Bonada N., Rieradevall M. and Prat N., Macroinvertebrate community structure and biological traits related to flow permanence in a Mediterranean river network, *Hydrobiologia* **589**, 2007, 91–106.
- Boulton A.J., Over-summering refuges of aquatic macroinvertebrates in two intermittent streams in central Victoria, *T. Roy. Soc. South. Aust.* **31**, 1989, 23–34.

- Boulton A.J., Parallels and contrasts in the effects of drought on stream macroinvertebrate assemblages, *Freshw. Biol.* **48**, 2003, 1173–1185.
- Boulton A.J., Sheldon F., Thoms M.C. and Stanley E.H., Problems and constraints in managing rivers with variable flow regimes, In: Boon P.J., Davies B.R. and Petts G.E., (Eds.), *Global Perspectives on River Conservation: Science, Policy and Practice*, 2000, John Wiley & Sons; London, 415–425.
- Brown C.A., Joubert A.R., Beuster J., Greyling A. and King J.M., DRIFT: DSS Software Development for Integrated Flow Assessments. WRC Report, 2013.
- Buffagni A., Armanini D.G. and Erba S., Does the lentic-lotic character of rivers affect invertebrate metrics used in the assessment of ecological quality?, *J. Limnol.* **68**, 2009, 92–105, <http://dx.doi.org/10.4081/jlimnol.2009.92>.
- Buytaert W., Zulkafli Z., Grainger S., Acosta L., Alemie T.C., Bastiaensen J., De Bièvre B., Bhusal J., Clark J., Dewulf A., Foggin M., Hannah D.M., Hergarten C., Isaeva A., Karpouzoglou T., Pandeya B., Paudel D., Sharma K., Steenhuis T., Tilahun S., Van Hecken G. and Zhumanova M., Citizen science in hydrology and water resources: opportunities for knowledge generation, ecosystem service management, and sustainable development, *Front. Earth Sci.* **2**, 2014, 1–21.
- CHJ, Jucar River Basin Management Plan: 2016–2021. Technical Report, 2016, Spanish Ministry of the Environment.
- Cid N., Verkaik I., García-Roger E.M., Rieradevall M., Bonada N., Sánchez-Montoya M.M., Gómez R., Suárez M.L., Vidal-Abarca M.R., Demartini D. and Buffagni A., A biological tool to assess flow connectivity in reference temporary streams from the Mediterranean Basin, *Sci. Total Environ.* **540**, 2016, 178–190.
- Cid N., Bonada N., Carlson S.M., Grantham T.E., Gasith A. and Resh V.H., High variability is a defining component of Mediterranean-climate rivers and their biota, *Water* **9**, 2017, 1–24.
- CIS, Ecological Flows in the Implementation of the Water Framework Directive. Common Implementation Strategy, Guidance Document No. 31. European Commission. Technical Report, 2015. <http://dx.doi.org/10.2779/775712>.
- Corti R. and Datry T., Terrestrial and aquatic invertebrates in the riverbed of an intermittent river: parallels and contrasts in community organisation, *Freshw. Biol.* **61**, 2015, 1308–1320.
- Custodio E., Andreu-Rodes J.M., Aragón R., Estrela T., Ferrer J., García-Aróstegui J.L., Manzano M., Rodríguez-Hernández L., Sahuquillo A. and del Villar A., Groundwater intensive use and mining in south-eastern peninsular Spain: hydrogeological, economic and social aspects, *Sci. Total Environ.* **559**, 2016, 302–316.
- Datry T., Larned S.T. and Tockner K., Intermittent rivers: a challenge for freshwater ecology, *Bioscience* **64**, 2014a, 229–235, <http://dx.doi.org/10.1093/biosci/bit027>.
- Datry T., Larned S.T., Fritz K.M., Bogan M.T., Wood P.J., Meyer E.I. and Santos A.N., Broad-scale patterns of invertebrate richness and community composition in temporary rivers: effects of flow intermittence, *Ecography* **37**, 2014b, 94–104.
- Datry T., Pella H., Leigh C., Bonada N. and Hugueny B., A landscape approach to advance intermittent river ecology, *Freshw. Biol.* **61**, 2016, 1200–1213.
- Davis J., Pavlova A., Thompson R. and Sunnucks P., Evolutionary refugia and ecological refuges: key concepts for conserving Australian arid zone freshwater biodiversity under climate change, *Glob. Chang. Biol.* **19**, 2013, 1970–84.
- De Girolamo A.M., Lo Porto A., Pappagallo G., Tzoraki O. and Gallart F., The hydrological status concept: application at a temporary river (Candelaro, Italy), *River Res. Appl.* **31**, 2015, 892–903.
- Dolédec S., Tilbian J. and Bonada N., Temporal variability in taxonomic and trait compositions of invertebrate assemblages in two climatic regions with contrasting flow regimes, *Sci. Total Environ.* **599**–600, 2017, 1912–1921.
- Döll P. and Schmied H.M., How is the impact of climate change on river flow regimes related to the impact on mean annual runoff? A global-scale analysis, *Environ. Res. Lett.* **7**, 2012, 14–37.
- European Commission, Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for community action in the field of water policy, Off. J. Eur. Communities 2000 <http://eur-lex.europa.eu/legal-content/en/TXT/?uri=CELEX:32000L0060>, accessed 05.11.16.
- European Commission, Communication from the Commission to the European Parliament the Council the European Economic and Social Committee and the Committee of the Regions. A Blueprint to Safeguard Europe's Water Resources, Brussels, <http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:52012DC0673>, 2012, accessed 17 Dec. 2016.
- Friberg N., Bonada N., Bradley D.C., Dunbar M.J., Edwards F.K., Grey J., Hayes R.B., Hildrew A.G., Lamoroux N., Trimmer M. and Woodward G., Biomonitoring of human impacts in freshwater ecosystems: the good, the bad and the ugly, *Adv. Ecol. Res.* **44**, 2011, 1–68.

- Fritz K.M., Johnson B.R. and Walters D.M., Field Operations Manual for Assessing the Hydrologic Permanence and Ecological Condition of Headwater Streams. EPA/600/R-06/126, 2006, US Environmental Protection Agency, Office of Research and Development; Washington DC <http://www.epa.gov/eerd/methods/headwater.html>, accessed 2 Sep. 2012.
- Gallart F., Prat N., Garca-Roger E.M., Latron J., Rieradevall M., Llorens P., Barbera G.G., Brito D., De Girolamo A.M., Lo Porto A., Buffagni A., Erba S., Neves R., Nikolaidis N.P., Perrin J.L., Querner E.P., Quinonero J.M., Tournoud M.G., Tzoraki O., Skoulidakis N., Gamez R., Gomez R. and Froebrich J., A novel approach to analysing the regimes of temporary streams in relation to their controls on the composition and structure of aquatic biota, *Hydrol. Earth Syst. Sci.* **16**, 2012, 3165–3182, <http://dx.doi.org/10.5194/hess-16-3165-2012>.
- Gallart F., Llorens P., Latron J., Cid N., Rieradevall M. and Prat N., Validating alternative methodologies to estimate the regime of temporary rivers when flow data are unavailable, *Sci. Total Environ.* **565**, 2016, 1001–1010.
- García-Roger E.M., Sánchez-Montoya M.M., Gómez R., Suárez M.L., Vidal-Abarca M.R., Latron J., Rieradevall M. and Prat N., Do seasonal changes in habitat features influence aquatic macroinvertebrate assemblages in perennial versus temporary Mediterranean streams?, *Aquat. Sci.* **73**, 2011, 567–579, <http://dx.doi.org/10.1007/s00027-011-0218-3>.
- Graham .D J., and Midgley N.G., Technical Communication-Graphical Representation of Particle Shape using Triangular Diagrams: An Excel Spreadsheet Method, *Earth Surf. Proc. Land.* **25**, 2000, 1473-1478.
- Hawkins C.P., Olson J.R. and Hill R.A., The reference condition: predicting benchmarks for ecological and water quality assessments, *J. N. Am. Benthol. Soc.* **29**, 2010, 312–343.
- International River Foundation, Brisbane Declaration, <http://riverfoundation.org.au/wp-content/uploads/2017/02/THE-BRISBANE-DECLARATION.pdf>, 2007, accessed 2 Mar. 2017.
- Jacobson P.M., Dixon D.A., Leggett W.C., Marcy B.C., Jr. and Massengill R.R., The Connecticut river ecological study (1965–1973) revisited: ecology of the lower Connecticut river 1973–2003, *American Fisheries Society Monograph* **9**, 2004.
- Kerezszy A., Gido K., Magalhaes M.F. and Skelton P.H., The biota of intermittent rivers and ephemeral streams: fishes, In: Datry T., Bonada N. and Boulton A.J., (Eds.), *Intermittent Rivers and Ephemeral Streams: Ecology and Management*, 2017, Elsevier Inc.; Cambridge, MA, (in press).
- Kirkby M.J., Gallart F., Kjeldsen T.R., Irvine B.J., Froebrich J., Porto A.L. and Girolamo A.D., Classifying low flow hydrological regimes at a regional scale, *Hydrol. Earth. Syst. Sc.* **15**, 2011, 3741–3750.
- Larned S.T., Datry T., Arscott D.B. and Tockner K., Emerging concepts in temporary-river ecology, *Freshw. Biol.* **5**, 2010, 717–738.
- Leigh C., Bonada N., Boulton A.J., Hugueney B., Larned S.T., Vander Vorste R. and Datry T., Invertebrate assemblage responses and the dual roles of resistance and resilience to drying in intermittent rivers, *Aquat. Sci.* **78**, 2016a, 291–301, <http://dx.doi.org/10.1007/s00027-015-0427-2>.
- Leigh C., Boulton A.J., Courtwright J.L., Fritz K., May C.L., Walker R.H. and Datry T., Ecological research and management of intermittent rivers: an historical review and future directions, *Freshw. Biol.* **61**, 2016b 1181–1199.
- Luthy R.G., Sedlak D.L., Plumlee M.H., Austin D. and Resh V.H., Wastewater-effluent-dominated streams as ecosystem-management tools in a drier climate, *Front. Ecol. Environ.* **13**, 2015, 477–485.
- Mackay S., Marsh N., Sheldon F. and Kennard M., Low-Flow Hydrological Classification of Australia, 2012, National Water Commission; Canberra http://content.webarchive.nla.gov.au/gov/wayback/20160615064923/http://archive.nwc.gov.au/_data/assets/pdf_file/0018/21807/Hydrological-classification.pdf, accessed 8 Feb. 2017.
- Martínez Santa-María C. and Fernández Yuste J.A., IAHRIS 2.2. Indicators of Hydrologic Alteration in Rivers. Methodological Reference Manual, 2010, Spanish Ministry of the Environment. Polytechnic University of Madrid http://www.ecogesfor.org/pdf/METH_REF_MANUAL_IAHRIS_v2_2.pdf, accessed 8 Feb. 2017.
- Mazor R.D., Stein E.D., Ode P.R. and Schiff K., Integrating intermittent streams into watershed assessments: applicability of an index of biotic integrity, *Freshw. Sci.* **33**, 2014, 459–474.
- Munné A. and Prat N., Effects of Mediterranean climate annual variability on stream biological quality assessment using macroinvertebrate communities, *Ecol. Indic.* **11**, 2011, 651–662.
- Mykrä H., Saarinen T., Tolkkinen M., McFarland B., Hämäläinen H., Martinmäki K. and Kløve B., Spatial and temporal variability of diatom and macroinvertebrate communities: how representative are ecological classifications within a river system?, *Ecol. Indic.* **18**, 2012, 208–217, <http://dx.doi.org/10.1016/j.ecolind.2011.11.007>.

- Nadeau T.L. and Rains M.C., Hydrological connectivity between headwater streams and downstream waters: how science can inform policy, *J. Am. Water Resour. Assoc.* **43**, 2007, 118–133, <http://dx.doi.org/10.1111/j.1752-1688.2007.00010.x>.
- Nikolaïdis N.P., Demetropoulou L., Froebrich J., Jacobs C., Gallart F., Prat N., Lo Porto A., Papadoulakis V., Campana C., Skoulikidis N., Davy T., Bidoglio G., Bouraoui F., Kirkby M.J., Tournoud M.G., Polesello S., González-Barberá G., Cooper D., Gomez R., Sanchez M.M. and De Girolamo A.M., Towards a sustainable management of Mediterranean river basins — policy recommendations on management aspects of temporary river basins, *Water Policy* **15** (5), 2013, 830–849, <http://dx.doi.org/10.2166/wp.2013.158>.
- Poff N.L., A hydrogeography of unregulated streams in the United States and an examination of scale-dependence in some hydrological descriptors, *Freshw. Biol.* **36**, 1996, 71–91.
- Poff N.L. and Ward J.V., Implications of streamflow variability for lotic community structure: a regional analysis of streamflow patterns, *Can. J. Fish. Aquat. Sci.* **46**, 1989, 1805–1817.
- Poff N.L., Richter B.D., Arthington A.H., Bunn S.E., Naiman R.J., Kendy E., Acreman M., Apse C., Bledsoe B.P., Freeman M.C. and Henriksen J., The ecological limits of hydrologic alteration (ELOHA): a new framework for developing regional environmental flow standards, *Freshw. Biol.* **55**, 2010, 147–170.
- Prat N., Gallart F., Von Schiller D., Polesello S., García-Roger E.M., Latron J., Rieradevall M., Llorens P., Barberá G.G., Brito D. and De Girolamo A.M., The mirage toolbox: an integrated assessment tool for temporary streams, *River Res. Appl.* **30**, 2014, 1318–1334.
- Reyjol Y., Argillier C., Bonne W., Borja A., Buijse A.D., Cardoso A.C., Daufresne M., Kernan M., Ferreira M.T., Poikane S., Prat N., Solheim A.L., Stroffek S., Usseglio-Polatera P., Villeneuve B. and van de Bund W., Assessing the ecological status in the context of the European water framework directive: where do we go now?, *Sci. Total Environ.* **497**, 2014, 332–344, <http://dx.doi.org/10.1016/j.scitotenv.2014.07.119>.
- Richter B.D., Baumgartner J.V., Powell J. and Braun D.P., A method for assessing hydrologic alteration within ecosystems, *Conserv. Biol.* **10**, 1996, 1163–1174.
- Riley S.P., Busted G.T., Kats L.B., Vandergon T.L., Lee L.F., Dagit R.G., Kerby J.L., Fisher R.N. and Sauvajot R.M., Effects of urbanization on the distribution and abundance of amphibians and invasive species in southern California streams, *Conserv. Biol.* **19** (6), 2005, 1894–1907.
- Robinson C., Uehlinger U. and Monaghan M., Effects of a multiyear experimental flood regime on macroinvertebrates downstream of a reservoir, *Aquat. Sci.* **65**, 2003, 210–222, <http://dx.doi.org/10.1007/s00027-003-0663-8>.
- Robson B.J., Chester E.T., Mitchell B.D. and Matthews T.G., Disturbance and the role of refuges in mediterranean climate streams, *Hydrobiologia* **719**, 2013, 77–91, <http://dx.doi.org/10.1007/s10750-012-1371-y>.
- Rossouw L., Avenant M.F., Seaman M.T., King J.M., Barker C.H., du Preez P.J., Pelser A.J., Roos J.C., van Staden J.J., van Tonder G.J. and Watson M., Environmental Water Requirements in Non-perennial Systems, Water Research Commission, WRC Report No: 1414/1/05, <http://www.wrc.org.za/KnowledgeHubDocuments/ResearchReports/1414.pdf>, 2005, Accessed 12 Ap. 2017.
- Sabater S., Timoner X., Borner G., De Wilde M., Stromberg J.C. and Stella J.C., The biota of intermittent rivers and ephemeral streams: algae and vascular plants, In: Datry T., Bonada N. and Boulton A.J., (Eds.), *Intermittent Rivers and Ephemeral Streams: Ecology and Management*, 2017, Elsevier Inc.; Cambridge, MA, (in press).
- Sánchez-Montoya M.M., Moleón M., Sánchez-Zapata J.A. and Tockner K., Dry riverbeds: corridors for terrestrial vertebrates, *Ecosphere* **7**, 2016, 1–10 [doi.wiley.com/10.1002/ecs2.1508](http://dx.doi.org/10.1002/ecs2.1508).
- Schriever T.A., Bogan M.T., Boersma K.S., Cañedo-Argüelles M., Jaeger K.L., Olden J.D. and Lytle D.A., Hydrology shapes taxonomic and functional structure of desert stream invertebrate communities, *Freshw. Sci.* **34**, 2015, 399–409.
- Seaman M., Watson M., Avenant M., King J., Joubert A., Barker C., Esterhuysen S., Graham D., Kemp M., Le Roux P. and Prucha B., DRIFT-ARID: a method for assessing environmental water requirements (EWRs) for nonperennial rivers, *Water SA* **42**, 2016, 356–367.
- Sheldon F., Incorporating natural variability into the assessment of ecological health in Australian dryland rivers, *Hydrobiologia* **552**, 2005, 45–56, <http://dx.doi.org/10.1007/s10750-005-1504-7>.
- Skoulikidis N.T., Sabater S., Datry T., Morais M.M., Buffagni A., Dörflinger G., Zogaris S., Sánchez-Montoya M.M., Bonada N., Kalogianni E., Rosado J., Vardakas L., De Girolamo A.M. and Tockner K., Non-perennial Mediterranean rivers in Europe: status, pressures, and challenges for research and management, *Sci. Total Environ.* **577**, 2017, 1–18, <http://dx.doi.org/10.1016/j.scitotenv.2016.10.147>.

Soil Survey Division Staff, Soil Survey Manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18, 1993.

Stubbington R., England J., Wood P.J. and Sefton C.E.M., Temporary streams in temperate zones: recognizing, monitoring and restoring transitional aquatic-terrestrial ecosystems, *WIREs Water* **2017**, 2017a, e1223<http://dx.doi.org/10.1002/wat2.1223>.

Stubbington R., Bogan M.T., Bonada N., Boulton A.J., Datry T., Leigh C. and Vander Vorste R., The biota of intermittent rivers and ephemeral streams: aquatic invertebrates, In: Datry T., Bonada N. and Boulton A.J., (Eds.), *Intermittent Rivers and Ephemeral Streams: Ecology and Management*, 2017b, Elsevier Inc.; Cambridge, MA, (in press).

Thoms M.C. and Sheldon F., An ecosystem approach for determining environmental water allocations in Australian dryland river systems: the role of geomorphology, *Geomorphology* **47**, 2002, 153–168, [http://dx.doi.org/10.1016/S0169-555X\(02\)00085-5](http://dx.doi.org/10.1016/S0169-555X(02)00085-5).

Tonkin J.D., Bogan M.T., Bonada N., Rios-Touma B. and Lytle D.A., Seasonality and predictability shape temporal species diversity, *Ecology* **2017**, Accepted Author Manuscript. 10.1002/ecy.1761.

Turner D.S. and Richter H.E., Wet/dry mapping: using citizen scientists to monitor the extent of perennial surface flow in dryland regions, *Environ. Manag.* **47**, 2011, 497–505.

Uys M.C. and O'Keeffe J.H., Simple words and fuzzy zones: Early directions for temporary river research in South Africa, *Environ. Manag.* **21**, 1997, 517–531, <http://dx.doi.org/10.1007/s002679900047>.

Appendix A: Graphs used in TREHS




Figure A1: Aquatic States Frequency Graph (Gallart et al. 2012)

Information provided (output)

- Relative frequency (%) of the aquatic states¹ throughout the year².

Application

- Visual assessment of temporal variability and seasonal predictability for the selection of the most appropriate sampling methods (e.g. if the degree of seasonality is high, traditional flow-phase biomonitoring methods can be used by adapting the sampling calendar to the period with the highest probability of finding lotic conditions).

¹Defined in Gallart et al. (2012) as transient sets of aquatic mesohabitats occurring in a stream reach at a particular moment depending on the hydrological conditions. These are: flood conditions (*Hyperrheic*), flow with all mesohabitats connected (*Eurheic*), sequence of pools connected by flowing water threads (*Oligorheic*), isolated pools (*Arheic*), absence of surface water with the wet alluvium (*Hyporheic*) and completely dry river bed and alluvium (*Edaphic*).

²When input data are from flow records or simulations, the temporal scale is represented on a monthly basis (a), and when they correspond to interviews, *in situ* or photographic observations, the temporal scale is represented on a seasonal basis (b).




Figure A.2: Temporary Regime Plot (Gallart et al. 2012)

Information provided (output)

- Flow permanence (Mf) and its variability.
- Seasonal predictability (Sd6) and its variability.
- MIRAGE flow regime classification³.

Application

- Quantitative assessment of the flow regime.
- Comparison of several sites and/or different sources of input data (e.g. flow records, flow simulations, interviews, observations).
- Visual estimation of flow regime alteration (i.e. hydrological status).

³The flow regime types defined by Gallart et al. (2012) –i.e. Perennial (P), Intermittent-pools (I-P), Intermittent-dry (I-P) and Ephemeral (E)- are shown in this plot for overall comparison. A more detailed classification is conducted in the Flow-Pool-Dry plot (FPD).




Figure A.3: Flow – Pool – Dry plot

Information provided (output)

- Flow permanence (Mf) and its uncertainty (resolution).
- Pool permanence (Mp) and its uncertainty (resolution).
- Dry channel permanence (Md) and its uncertainty (resolution).
- TREHS temporary regime classification⁴.

Application

- Information on the presence of surface water when flow ceases (i.e. isolated pools).
- Comparison of several sites and/or different sources of input data (e.g. flow records, flow simulations, interviews, observations).
- Visual estimate of flow regime alteration (i.e. hydrological status).

⁴ The flow regime types defined in this paper are: Perennial (Pe), Quasi-perennial (Qp), Fluent-Stagnant (FS), Alternate-Fluent (AF), Stagnant (St), Alternate-Stagnant (AS), Alternate (Al), Occasional (Oc), Episodic (Ep). See Table 1 and Figure 2 for more detailed information.